arXiv:1006.2033v1 [math.NT] 10 Jun 2010

SOME IDENTITIES ON THE q-BERNSTEIN POLYNOMIALS, q-STIRLING NUMBERS AND q-BERNOULLI NUMBERS

TAEKYUN KIM, JONGSUNG CHOI, AND YOUNG-HEE KIM

Abstract In this paper, we consider the q-Bernstein polynomials on $\mathbb{Z}_p$ and investigate some interesting properties of q-Bernstein polynomials related to q-Stirling numbers and Carlitz's type q-Bernoulli numbers.

1. Introduction

Let $C[0,1]$ denote the set of continuous functions on $[0,1]$. Then Berstein operator for $f \in C[0,1]$ is defined as

$$\mathbb{B}_n(f)(x) = \sum_{k=0}^{n} f(\frac{k}{n})\binom{n}{k}x^k(1-x)^{n-k} = \sum_{k=0}^{n} f(\frac{k}{n})B_{k,n}(x),$$

for $k, n \in \mathbb{Z}_+$, where $B_{k,n}(x) = \binom{n}{k}x^k(1-x)^{n-k}$ is called the Bernstein polynomial of degree n (see [1, 9, 10]).

In [9], Phillips introduced the q-extension of Bernstein polynomials and Kim-Jang-Yi proposed the modified q-Bernstein polynomials of degree n, which are different q-Berstein polynomials of Phillips (see [1]).

Let q be regarded as either a complex number $q \in \mathbb{C}$ or a p-adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, then we always assume that $|q| < 1$. If $q \in \mathbb{C}_p$, we normally assume that $|1-q|_p < p^{-\frac{1}{p-1}}$, which yields the relation $q^x = \exp(x \log q)$ for $|x|_p \le 1$ (see [1-11]).

Here, the symbol $|\cdot|_p$ stands for the p-adic absolute on $\mathbb{C}_p$ with $|p|_p = 1/p$. The q-bosonic natural numbers are defined by $[n]_q = \frac{1-q^n}{1-q} = 1 + q + \cdots + q^{n-1}$ $(n \in \mathbb{N})$, and the q-factorial is defined by $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$. In this paper we use the notation for the Gaussian binomial coefficients in the form of

$$\binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q![k]_q!} = \frac{[n]_q[n-1]_q \cdots [n-k+1]_q}{[k]_q!}, \quad \text{(see [3, 8])}.$$

Note that $\lim_{q\to 1}\binom{n}{k}_q = \binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{n!}$.

Let p be a fixed prime number. Throughout this paper, the symbol $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ denote the ring of p-adic integers, the field of p-adic rational numbers, the complex number field, and the completion of algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$.

2000 Mathematics Subject Classification : 11B68, 11B73, 41A30, 05A30, 65D15.

Key words and phrases : Bernstein polynomial, Bernstein operator, Bernoulli numbers and polynomials, Stirling number.

Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable function on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the p-adic q-integral on $\mathbb{Z}_p$ is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \quad \text{(see [4, 5, 6])}.$$

Carlitz' q-Bernoulli number $\beta_{k,q}$ can be defined respectively by $\beta_{0,q} = 1$ and by the rule that $q(q\beta+1)^k - \beta_{k,q}$ is equal to 1 if $k = 1$ and to 0 if $k > 1$ with the usual convention of replacing β^i by $\beta_{i,q}$ (see [5]). As was shown in [5], Carlitz's q-Bernoulli numbers can be represented by p-adic q-integral on $\mathbb{Z}_p$ as follows :

$$\int_{\mathbb{Z}_p} [x]_q^n d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} [x]_q^n q^x = \beta_{n,q}, \quad n \in \mathbb{Z}_+. \tag{1}$$

The k-th order factorial of the q-number $[x]_q$, which is defined by

$$[x]_{k,q} = [x]_q [x-1]_q \cdots [x-k+1]_q = \frac{(1-q^x)(1-q^{x-1})\cdots(1-q^{x-k+1})}{(1-q)^k},$$

is called the q-factorial of x of order k (see [3]). Thus, we note that $\binom{x}{k}_q = \frac{[x]_{k,q}}{[k]_q!}$.

In this paper, we consider q-Bernstein polynomials on $\mathbb{Z}_p$ and we investigate some interesting properties of q-Bernstein polynomials related q-Stirling numbers and Carlitz's q-Bernoulli numbers.

2. q-BERNSTEIN POLYNOMIALS RELATED TO q-STIRLING NUMBERS AND q-BERNOULLI NUMBERS

In this section, we assume that $q \in \mathbb{C}_p$ with $|1-q|_p < p^{-\frac{1}{p-1}}$. For $f \in UD(\mathbb{Z}_p)$, we consider q-Bernstein type operator on $\mathbb{Z}_p$ as follows :

$$\begin{aligned} \mathbb{B}_{n,q}(f)(x) &= \sum_{k=0}^{n} f(\frac{k}{n}) \binom{n}{k} [x]_q^k [1-x]_q^{n-k} \\ &= \sum_{k=0}^{n} f(\frac{k}{n}) B_{k,n}(x,q), \end{aligned} \tag{2}$$

for $k, n \in \mathbb{Z}_+$, where $B_{k,n}(x,q) = \binom{n}{k}[x]_q^k[1-x]_q^{n-k}$ is called q-Bernstein type polynomials of degree n (see [1]).

Let $(Eh)(x) = h(x+1)$ be the shift operator. Consider the q-difference operator as follows :

$$\Delta_q^n = \prod_{i=1}^{n} (E - q^{i-1} I), \tag{3}$$

where $(Ih)(x) = h(x)$. From (3), we note that

$$f(x) = \sum_{n\geq 0} \binom{x}{n}_q \Delta_q^n f(0), \quad \text{(see [3])}, \tag{4}$$

where

$$\Delta_q^n f(0) = \sum_{k=0}^{n} \binom{n}{k}_q (-1)^k q^{\binom{k}{2}} f(n-k). \tag{5}$$

The q-Stiring number of the first kind is defined by

$$\prod_{k=1}^{n}(1+[k]_q z)=\sum_{k=0}^{n} S_1(n,k:q)z^k, \tag{6}$$

and the q-Stiring number of the second kind is also defined by

$$\prod_{k=1}^{n}(\frac{1}{1+[k]_q z})=\sum_{k=0}^{n} S_2(n,k:q)z^k, \quad \text{(see [3]).} \tag{7}$$

By (3), (4), (5), (6) and (7), we see that

$$S_2(n,k:q)z^k=\frac{q^{-\binom{k}{2}}}{[k]_q!}\sum_{j=0}^{k}(-1)^j q^{\binom{j}{2}}\binom{k}{j}_q [k-j]_q^n=\frac{q^{-\binom{k}{2}}}{[k]_q!}\Delta_q^k 0^n, \quad \text{(see [1]).} \tag{8}$$

For $q\in\mathbb{C}_p$ with $|1-q|_p<p^{-\frac{1}{p-1}}$, we have that for $k,n\in\mathbb{Z}_+$,

$$\begin{aligned} F_q^{(k)}(t,x) &= \frac{t^k e^{[1-x]_q t}[x]_q^k}{k!}=[x]_q^k\sum_{n=0}^{\infty}\binom{n+k}{k}[1-x]_q^n\frac{t^{n+k}}{(n+k)!} \\ &= \sum_{n=k}^{\infty}\binom{n}{k}[x]_q^k[1-x]_q^{n-k}\frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty}B_{k,n}(x,q)\frac{t^n}{n!}, \quad \text{(see [1]).} \end{aligned}$$

Thus, we note that $\frac{t^k e^{[1-x]_q t}[x]_q^k}{k!}$ is the generating function of q-Berstein polynomials (see [1]). It is easy to show that

$$[1-x]_q^{n-k}=\sum_{m=0}^{\infty}\sum_{l=0}^{n-k}\binom{l+m-1}{m}\binom{n-k}{l}(-1)^{l+m}q^l[x]_q^{l+m}(q-1)^m. \tag{9}$$

By (1), (2) and (9), we obtain the following theorem.

Theorem 1. *For $k,n\in\mathbb{Z}_+$ with $n\geq k$, we have*

$$\begin{aligned} &\int_{\mathbb{Z}_p} B_{k,n}(x,q)d\mu_q(x) \\ &\quad =\sum_{m=0}^{\infty}\sum_{l=0}^{n-k}\binom{l+m-1}{m}\binom{n-k}{l}(-1)^{l+m}q^l(q-1)^m\beta_{l+m+k,q}, \end{aligned}$$

where $\beta_{n,q}$ are the n-th Carlitz q-Bernoulli numbers.

In [3], it is known that

$$[x]_q^n=\sum_{k=0}^{n}q^{\binom{k}{2}}\binom{x}{k}_q[k]_q!S_2(n,k:q), \tag{10}$$

and

$$\frac{\sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}B_{k,n}(x,q)}{([x]_q+[1-x]_q)^{n-i}}=[x]_q^i, \quad \text{for } i\in\mathbb{N}, \quad \text{(see [1]).} \tag{11}$$

By (2) and (11), we see that for $i \in \mathbb{N}$,

$$\begin{aligned}[x]_q^i = &\sum_{m=0}^{\infty}\sum_{k=i-1}^{n}\sum_{l=0}^{m+n-k}\sum_{p=0}^{\infty}\frac{\binom{k}{i}\binom{n}{k}}{\binom{n}{i}}\binom{l+p-1}{p} \\ &\times\binom{m+n-k}{l}\binom{n-i+m-1}{m}(-1)^{l+p+m}q^l(q-1)^p[x]_q^{n-i-m+k+p+l}.\end{aligned} \tag{12}$$

By (12), we obtain the following theorem.

Theorem 2. *For $k, n \in \mathbb{Z}_+$ and $i \in \mathbb{N}$, we have*

$$\begin{aligned}\beta_{i,q} = &\sum_{m=0}^{\infty}\sum_{k=i-1}^{n}\sum_{l=0}^{m+n-k}\sum_{p=0}^{\infty}\frac{\binom{k}{i}\binom{n}{k}}{\binom{n}{i}}\binom{l+p-1}{p} \\ &\times\binom{m+n-k}{l}\binom{n-i+m-1}{m}(-1)^{l+p+m}q^l(q-1)^p\beta_{n-i-m+k+p+l,q}.\end{aligned}$$

From (10) and (11), we note that

$$\frac{\sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}B_{k,n}(x,q)}{([x]_q+[1-x]_q)^{n-i}} = \sum_{k=0}^{i} q^{\binom{k}{2}}\binom{x}{k}_q [k]_q! S_2(i,k:q). \tag{13}$$

In (3), it is known that

$$\int_{\mathbb{Z}_p}\binom{x}{n}_q d\mu_q(x) = \frac{(-1)^n}{[n+1]_q}q^{(n+1)-\binom{n+1}{2}}. \tag{14}$$

By (13), (14) and Theorem 2, we have

$$\beta_{n,q} = q\sum_{k=0}^{m}\frac{[k]_q!}{[k+1]_q}(-1)^k S_2(k, n-k, : q).$$

For $S_2(n,k:q)$, we see that

$$S_2(n,k:q) = \frac{1}{(1-q)^k}\sum_{j=0}^{k}(-1)^{k-j}\binom{k+n}{k-j}\binom{j+n}{j}_q, \quad \text{(see [3])}, \tag{15}$$

and

$$\binom{n}{k}_q = \sum_{j=0}^{n}\binom{n}{j}(q-1)^{j-k}S_2(k, j-k:q).$$

By simple calculation, we show that

$$\begin{aligned} q^{nx} &= \sum_{k=0}^{n}(q-1)^k q^{\binom{k}{2}}\binom{n}{k}_q [x]_{k,q} \\ &= \sum_{m=0}^{n}\{\sum_{k=m}^{n}(q-1)^k\binom{n}{k}_q S_1(k,m:q)\}[x]_q^m, \end{aligned} \tag{16}$$

and

$$\int_{\mathbb{Z}_p} q^{nx}d\mu_q(x) = \sum_{m=0}^{\infty}\binom{n}{m}(q-1)^m\beta_{m,q}. \tag{17}$$

By (16) and (17), we see that

$$\binom{n}{m}_q = \sum_{k=m}^{n} (q-1)^{-m+k} \binom{n}{k}_q S_1(k, m : q).$$

Therefore, we obtain the following theorem.

Theorem 3. *For $k, n \in \mathbb{Z}_+$, we have*

$$B_{k,n}(x, q) = \sum_{m=k}^{n} (q-1)^{-k+m} \binom{n}{m}_q S_1(m, k : q)[x]_q^k [1-x]_q^{n-k}.$$

From the definition of the q-Stirling numbers of the first kind, we derive

$$q^{\binom{n}{2}} \binom{x}{n}_q [n]_q! = [x]_{n,q} q^{\binom{n}{2}} = \sum_{k=0}^{n} S_1(n, k : q)[x]_q^k. \tag{18}$$

By (13) and (18), we obtain the following theorem.

Theorem 4. *For $k, n \in \mathbb{Z}_+$ and $i \in \mathbb{N}$, we have*

$$\frac{\sum_{k=i-1}^{n} \frac{\binom{k}{i}}{\binom{n}{i}} B_{k,n}(x, q)}{([1-x]_q + [x]_q)^{n-i}} = \sum_{k=0}^{i} (\sum_{l=0}^{k} S_1(n, l : q) S_2(i, k : q)[x]_q^l).$$

By Theorem 2 and Theorem 4, we obtain the following result.

Corollary 5. *For $i \in \mathbb{N}$, we have*

$$\beta_{i,q} = \sum_{k=0}^{i} (\sum_{l=0}^{k} S_1(n, l : q) S_2(i, k : q) \beta_{l,q}),$$

where $\beta_{i,q}$ are the i-th Carlitz q-Bernoulli numbers.

In [3], the q-Bernoulli polynomials of order $k\,(\in \mathbb{Z}_+)$ are defined by

$$\beta_{n,q}^{(k)}(x) = \frac{1}{(1-q)^n} \sum_{i=0}^{n} (-1)^i \binom{n}{i} q^{ix} \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k-\text{times}} q^{\sum_{l=1}^{k}(k-l+i)x_l} d\mu_q(x_1) \cdots d\mu_q(x_k). \tag{19}$$

From (19), we note that

$$\beta_{n,q}^{(k)} = \frac{1}{(1-q)^n} \sum_{i=0}^{n} (-1)^i \binom{n}{i} \frac{(i+k) \cdots (i+1)}{[i+k]_q \cdots [i+1]_q} q^{ix}, \quad \text{(see [3])}.$$

The inverse q-Bernoulli polynomial of order k are also defined by

$$\beta_{n,q}^{(-k)}(x) = \frac{1}{(1-q)^n} \sum_{i=0}^{n} \frac{(-1)^n \binom{n}{i} q^{xi}}{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} q^{\sum_{l=1}^{k}(k-l+i)x_l} d\mu_q(x_1) \cdots d\mu_q(x_k)}, \tag{20}$$

(see [3]). In the special case $x = 0$, $\beta_{n,q}^{(k)} = \beta_{n,q}^{(k)}(0)$ are called the n-th q-Bernoulli numbers of order k and $\beta_{n,q}^{(-k)} = \beta_{n,q}^{(-k)}(0)$ are called the n-th inverse q-Bernoulli numbers of order k.

By (20), we see that

$$\begin{aligned}\beta_{k,q}^{(-n)} &= \frac{1}{(1-q)^k}\sum_{j=0}^{k}(-1)^j\binom{k}{j}\frac{[j+n]_q\cdots[j+1]_q}{(j+n)\cdots(j+1)} \\ &= \frac{1}{(1-q)^k}\sum_{j=0}^{k}(-1)^j\frac{\binom{k}{j}}{(j+n)\cdots(j+1)}[n]_q!\binom{j+n}{n}_q \qquad (21) \\ &= \frac{1}{(1-q)^k}\sum_{j=0}^{k}(-1)^j\frac{\binom{k+n}{n-j}}{\binom{k+n}{n}}\binom{j+n}{n}_q\frac{[n]_q!}{n!} \\ &= \frac{[n]_q!}{\binom{k+n}{n}n!}\{\frac{1}{(1-q)^k}\sum_{j=0}^{k}(-1)^j\binom{k+n}{k-j}\binom{j+n}{n}_q\}.\end{aligned}$$

From (15) and (21), we note that

$$S_2(n,k:q)=\binom{k+n}{n}\frac{[n]_q!}{n!}\beta_{k,q}^{(-n)}. \qquad (22)$$

By (13) and (22), we obtain the following theorem.

Theorem 6. *For $k,n\in\mathbb{Z}_+$ and $i\in\mathbb{N}$, we have*

$$\frac{\sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}B_{k,n}(x,q)}{([1-x]_q+[x]_q)^{n-i}}=\sum_{k=0}^{i}q^{\binom{k}{2}}\binom{x}{k}_q[k]_q!\binom{k+i}{i}\frac{[i]_q!}{i!}\beta_{k,q}^{(-i)}.$$

It is not difficult to show that

$$\begin{aligned}q^{\binom{n}{2}}\binom{x}{n}_q &= \frac{1}{[n]_q!}\prod_{k=0}^{n-1}([x]_q-[k]_q) \qquad (23) \\ &= \frac{1}{[n]_q!}\sum_{k=0}^{n}(-1)^k[x]_q^{n-k}S_1(n-1,k:q). \qquad (24)\end{aligned}$$

By Theorem 4 and (23), we obtain the following result.

Corollary 7. *For $k,n\in\mathbb{Z}_+$ and $i\in\mathbb{N}$, we have*

$$\frac{\sum_{k=i-1}^{n}\frac{\binom{k}{i}}{\binom{n}{i}}B_{k,n}(x,q)}{([1-x]_q+[x]_q)^{n-i}}=\sum_{k=0}^{i}\sum_{j=0}^{k}(-1)^j[x]_q^{n-j}S_1(k-1,j:q)\binom{k+i}{i}\frac{[i]_q!}{i!}\beta_{k,q}^{(-i)}.$$

REFERENCES

[1] T. Kim, L.-C. Jang, H. Yi, *Note on the modified q-Berstein polynomials*, arXiv 1005.4293.

[2] T. Kim, *q-Volkenborn integration* , Russ. J. Math. Phys. **9** (2002), no. 3, 288–299.

[3] T. Kim, *q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), 51–57.

[4] T. Kim, *Note on the Euler q-zeta functions*, J. Number Theory **129** (2009), 1798–1804.

[5] T. Kim, *On a q-analogue of the p-adic log gamma functions and related integrals*, J. Number Theory **76** (1999), 320–329.

[6] T. Kim, *Barnes-type multiple q-zeta functions and q-Euler polynomials*, J. Physics A : Math. Theor., **43** (2010), 255201.

[7] T. Kim, *Power series and asymptotic series associated with the q-analog of the two-variable p-adic L-function*, Russ. J. Math. Phys. **12** (2005), 186–196.

[8] T. Kim, *On p-adic interpolating function for q-Euler numbers and its dericatives*, J. Math. Anal. Appl. **339** (2008), 598–608.

[9] G. M. Phillips, *Bernstein polynomials based on the q-integer*, Annals of Numerical Analysis **4** (1997), 511–518.

[10] S. Bernstein, *Demonstration du theoreme de Weierstrass, fondee sur le calcul des probabilities*, Commun. Soc. Math. Kharkow (2) **13** (1912-1913), 1–2.

[11] N. K. Govil, V. Gupta, *Convergence of q-Meyer-Konig-Zeller-Purrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 181–189.

Taekyun Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `tkkim@kw.ac.kr`

Jongsung Choi. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `jeschoi@kw.ac.kr`

Young-Hee Kim. Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea,
E-mail address: `yhkim@kw.ac.kr`